

- 1. This document was created to support maximum accessibility for all learners. If you would like to print a hard copy of this document, please follow the general instructions below to print multiple slides on a single page or in black and white.
- 2. If you are viewing this course as a recorded course after the live webinar, you can use the scroll bar at the bottom of the player window to pause and navigate the course.
- 3. This handout is for reference only. Non-essential images have been removed for your convenience. Any links included in the handout are current at the time of the live webinar, but are subject to change and may not be current at a later date.
- 4. Copyright: Images used in this course are used in compliance with copyright laws and where required, permission has been secured to use the images in this course. All use of these images outside of this course may be in violation of copyright laws and is strictly prohibited.

How to print Handouts

- On a PC
 - Open PDF
 - Click Print
 - Choose # of pages per sheet from dropdown menu
 - Choose Black and White from "Color" dropdown
- On a Mac
 - Open PDF in Preview
 - Click File
 - Click Print
 - Click dropdown menu on the right "preview"
 - Click layout
- Choose # of pages per sheet from dropdown menu
- Checkmark Black & White if wanted.
- If more details needed please visit our FAQ page: https://www.speechpathology.com/help



No part of the materials available through the continued.com site may be copied, photocopied, reproduced, translated or reduced to any electronic medium or machine-readable form, in whole or in part, without prior written consent of continued.com, LLC. Any other reproduction in any form without such written permission is prohibited. All materials contained on this site are protected by United States copyright law and may not be reproduced, distributed, transmitted, displayed, published or broadcast without the prior written permission of continued.com, LLC. Users must not access or use for any commercial purposes any part of the site or any services or materials available through the site.

Technical issues with the Recording?

- Clear browser cache using these instructions
- Switch to another browser
- Use a hardwired Internet connection
- Restart your computer/device

Still having issues?

- Call 800-242-5183 (M-F, 8 AM-8 PM ET)
- Email customerservice@speechpathology.com



TBI MASTER COURSE #3

Traumatic Brain Injury: Outpatient and Return to Work – Case Study Applications

Janice Osborne-Dowdy, MS, CCC/SLP, CBIS Speech-Language Pathologist NeuroRestorative. Ashland, KY

Steven Wheeler, PhD, OTR/L, FAOTA, CBIS Professor, Occupational Therapy West Virginia University



Janice Dowdy, MS, CCC-SLP, CBIS

Janice Osborne Dowdy, MS/CCC-SLP, CBIS, is the lead speech-language pathologist at NeuroRestorative Ashland, Kentucky. Janice is a Certified Brain Injury Specialist, through the Academy of Certified Brain Injury Specialists. She is a graduate of Eastern Kentucky University and East Tennessee State University. Janice has been a practicing SLP for 19 years treating individuals with cognitive/communication/swallowing deficits as the result of CVA, brain injury, head and neck cancer, progressive neurological diseases, as well as other disease processes. She has worked in acute care, long term acute care, skilled nursing, and acute rehabilitation in inpatient, outpatient, and day treatment programs both as a staff therapist, and in various management roles prior to coming to NeuroRestorative. Janice also worked for Chattanooga Group/EMPI as Clinical Specialist for Vital Stim Therapy, traveling the USA providing education and consulting services to physicians and therapists in the management and treatment of dysphagia utilizing the modality of neuromuscular electrical stimulation (NMES). Janice has worked in the Ashland program since 2013.





Steven Wheeler, PhD, OTR/L, CBIS

Dr. Steven Wheeler is a professor and occupational therapy division chairperson at the West Virginia University School of Medicine. Prior to this, he served in a similar role at the University of Cincinnati. Dr. Wheeler is also principal investigator on the West Virginia TBI State Project which seeks to monitor, expand, and improve the lives of TBI survivors and caregivers through education, advocacy, and community outreach. Dr. Wheeler received his PhD in Health-Related Sciences with a Specialization in Occupational Therapy from Virginia Commonwealth University. He has conducted TBI related presentations nationally and internationally over the past 20 years and has numerous publications on the topic including co-authoring the American Occupational Therapy Association's "Occupational Therapy Practice Guidelines for Adults with Traumatic Brain Injury."



continued

- Presenter Disclosure: Financial: Steven Wheeler and Janice Dowdy have received an honorarium for presenting this course. Nonfinancial: Steven Wheeler and Janice Dowdy have no relevant non-financial relationships to disclose.
- Content Disclosure: This learning event does not focus exclusively on any specific product or service.
- Sponsor Disclosure: This course is presented by OccupationalTherapy.com.



Learning Outcomes

After this course, participants will be able to:

- 1. Recognize common cognitive and perceptual impairments associated with traumatic brain injury.
- 2. Recognize the impact of cognitive and perceptual deficits on the performance of daily activities.
- 3. Identify approaches to assessment and intervention in the outpatient and community-based setting.



The Challenge of TBI Rehabilitation

Getting from here



To here






The Complexity of TBI and Human Performance

- Effective TBI intervention is a combination of:
 - Understanding TBI
 - Understanding your patient/client
 - Understanding activity and activity analysis
 - When an individual's needs are considered in the context of instructions or expectations, the learning/ rehabilitation process is accelerated.



Case Background - Drew





Step 1:

Accurate assessment of impairments is critical to successful neurological rehabilitation

But first ...



Assessment Foundations – Taking a "Top Down" Approach

- ClientCenteredness
- Rapport Building
- Establishing the Basis for Contextualized Rehabilitation





Rehabilitation "On the Average"

Decontextualized Approach Contextualized Approach Interview Set goals based on **PRIORITIZE** patient+family to norms/societal **ADLs/IADLs** identify and assumptions and **EVALUATE** prioritize goals where performance is breaking down Patients with increased portions of time in contextualized therapy have better outcomes Direct work on ADL/ **IADL** through **Develop strategies to** Strengthen motor, shaping, chaining, FACILITATE PERFORMANCE perceptual, and/or gradual removal of cognitive of ADL/IADL supports, components of the compensatory ADL/IADL strategies, etc.



Findings Regarding Contextualized Treatment (Bogner et al., 2019)

- Increasing the proportion of treatment using a contextualized approach results in better community participation one year later
 - Patients with a 30% greater proportion of contextualized treatment were more likely to be out of the house 1-2 more days a week one year later.
- Patients with greater disability experienced more benefit in regard to selfcare and mobility than patients with less severe disability
- The findings do not indicate that decontextualized should not be used, but to use contextualized treatment whenever possible given the therapy goal.



Assessment of Performance – Where do I Start?

- By knowing where your patient and family wants to go, you can evaluate to determine the discrepancy between now and your target goal
- Standardized
 assessments and
 observation of task
 participation can both
 be beneficial.



Goals: A. Work B. School C. Relationship



Recovery from TBI: Considerations in Work and School Capacity

Cognition / Perception	Physical / Motor	Behavioral / Emotional
Orientation Attention and Concentration Memory Problem solving Flexible thinking Initiation Disinhibition Self-awareness Goal-setting	Balance Fine and gross motor skills Spasticity Ataxia Pain / Headache Endurance Sensory changes Speech Swallowing Seizure disorder Sleep disturbance	Impulsivity Apathy Depression Anxiety Personality changes Emotional lability Flat affect Social skills Mood swings Aggression



About TBI: Common Problems Processing and Understanding Information

- Taking a longer time to grasp what others are saying
- Taking more time to understand and follow directions
- Have trouble following the plot of television shows or movies
- Taking longer to read and understand information such as a newspaper or book
- Slower to complete even routine tasks such as getting dressed or cooking a meal




About TBI: Common Problems Learning and Remembering New Information

- Related to memory deficits
- Difficulty with learning new routines and systems
- May remember some parts of events or conversations, but may appear to confabulate parts to "fill in the gap"



Establishing Competencies in Relation to Goals






- A. Cognition
- **B. Self-Awareness**
- C. Perception
- D. Communication

Dynamic Lowenstein Occupational Therapy Cognitive Assessment



Establishing Baselines Relative to Goals




How would we describe performance?

Dynamic Lowenstein Occupational Therapy Cognitive Assessment



Establishing Baselines Relative to Goals



Beery Test of Visual-Motor Integration (VMI)



Reading, writing, keyboard skills



Process of Storing Memory

- Attention
- Encoding Information
 - Being able to "understand" the information
- Storage
 - Moving information to a specific area of the brain
- Retrieval
 - Searching the "files" and pulling out a memory

continued


Establishing Physical Capacity Related to Work and School

- Baseline Assessment
 - Lifting
 - Floor-knuckle
 - Knuckle to shoulder
 - Shoulder to overhead
 - Carrying
 - Fine Motor
 - Endurance / Positional Tolerances
 - Pain





Return to Work Progression







Return to Work Progression

- Work capacity assessment
- Clinic based simulated work activities
- Remediation / compensation of impairments
- Volunteer activities
- Paid work

*** Requires establishing partnerships / advocacy



Making a Difference in the Clinic and Community

- By understanding the challenges experienced during recovery, therapists can make a big difference in the lives of clients and families
- Increasing participation in meaningful roles and pleasant activity (Bombardier, et al., 2017) and aerobic exercise (Weinstein et al., 2017) positively impacts mood





Important Considerations: Language and Communication Problems Post TBI

- Difficulty thinking of the correct word
- Trouble starting, following, understanding conversation
- Difficulty staying on topic
- Difficulty expressing thoughts in an organized manner
- Difficulty communicating thoughts and feelings using facial expressions and tone of voice
- Having trouble reading others thoughts and feelings
- Misunderstanding jokes and sarcasm being very "black and white"



Important Considerations: Planning and Organization Problems after TBI

- Difficulty planning schedules and following them
- Trouble with things requiring multiple steps from simple to complex tasks
- May lose or misplace items frequently, blame others for moving items or taking from them



Important Considerations: Problems with Reasoning, Problem Solving, and Judgment

- Difficulty realizing there is even a problem to start with...
 - Rigid thinking trouble analyzing options and solutions or changing the way they think
 - Quick judgments
 - Acting without thinking about the consequences



Important Considerations: Inappropriate, Embarrassing, or Impulsive Behavior

- Denying cognitive problems
- Saying hurtful or insensitive things
- Cursing or acting "out of character" compared to pre-injury self
- Lacking awareness of social boundaries
 - Sexually inappropriate
 - Invading personal space
 - Oversharing information with "strangers"



So can we "fix" any of these problems, and if so, how????



Remediation Vs. Compensation

- Two approaches
 - Fixing the problem Remediation
 - Coming up with ways to do the things you need to do, in a new way – Compensation
- Most progress and increase in skills can be seen from combination of the 2 approaches.



• Neuroplasticity: The brain's ability to reorganize itself by forming new neural connections throughout life. Neuroplasticity allows the neurons (nerve cells) in the brain to compensate for injury and disease and to adjust their activities in response to new situations or to changes in their environment.



- Use it or lose it
 - Not using functions of your brain can lead to no longer being able to use them
- Use it and improve it
 - Training a specific brain function can lead to an improvement in that function
- Specificity
 - The nature of the training dictates the nature of the plasticity
- Repetition Matters
 - Induction of plasticity requires sufficient repetition



- Intensity matters
 - Induction of plasticity requires sufficient training intensity
- Time matters
 - Different forms of plasticity occur at different times during the training
- Salience matters
 - The training experience must be sufficiently salient to induce plasticity
- Age matters
 - Training induced plasticity occurs more readily in young brains

(Kleim & Jones, 2008)



- Transference
 - Plasticity in response to one's training experience can enhance the acquisition of similar behaviors
- Interference
 - Plasticity in response to one experience can interfere with the acquisition of other behaviors.

(Kleim & Jones, 2008)




Treatment

- Individualize the educational information provided about the client's specific memory and cognitive difficulties
- Involve the client in setting specific goals and selecting memory aids
- Create opportunities for accurate self-monitoring
 - Providing a delay between a prediction for a memory task, allows cueing for memory aid, thus giving self-monitoring and self-control an explicit connection



Engaging the Client





Treatment

- Anosagnosia
 - Lack of awareness into deficits
- Anosagnosia is one of the biggest challenges in success with individuals with a TBI
 - Provide "safe" opportunities for failure
 - Example would be allowing the person to make a meal "their way" and it not turn out – burnt toast, runny brownies, yucky coffee
 - Breakdown and discuss task failures immediately after they occur – provide specific feedback.



Treatment Approaches

- Drill and Practice
- Dual Task Training
- Errorless learning
- Role playing
- Situational coaching



Case Example - Drill and Practice





Case Example – Social Discourse with Feedback





Life Skills Training / Coaching

 Teaching in specific life / job skills can facilitate community reintegration and provide valuable feedback to promote selfawareness

 Peer mentoring models have also been found to be effective







Compensatory Training – Examples

- Internal Memory Aids
 - Visual imagery
 - Category grouping
 - Mnemonics
 - Rehearsal strategies
- External Memory Aids
 - Written planning systems (calendar, notebook, planner appointment book)
 - Electronic planners (PDA, Blackberry, notepad on cellphone)
 - Computer based planning (Outlook, Google calendar) that links between devices




Compensation







Case Example – Compensation







Compensatory Training

- External Memory Aids
 - Items that can be task specific
 - Lists to check off and complete ADLS showering, doing laundry, grocery list
 - Post It notes (can easily get overwhelming without a "system")
 - Pill organizer
 - Key finder
 - Dry erase board/bulletin board
 - Television Assisted Prompting (TAP)
 - NeuroPage system



Compensatory Training

- Establishing both internal and external SUCCESSFUL memory aids requires them to be individualized to the client and re-evaluated as needed to modify and advance the complexity
 - Start at level of establishing orientation
 - Advance to higher level tasks cautious to establish success at each level



- Minimize distractions work in a quiet room
- Focus on one activity at a time
- Take a break when you feel tired
- Allow extra time to complete tasks
- Re-read and discuss new information, try to put it in "your own words"
- Try not to talk down to the person having difficulties, use a calm voice, and kind words
- Do not speak too fast or say too much at once



- Put together a structured routine of daily tasks and activities – and follow it – as close as possible, including sticking to planned times to eat, practice for therapy, complete basic ADLs, wake up, and go to bed.
- Set specific locations for keeping things a place for medication box, a hook to keep keys by door, table for purse/memory book/ items you need to take to therapy or work, keep a notebook next to the telephone for messages/details of conversation



- Schedule out the whole month Examples:
 - Grocery on Tuesday
 - Pay bills on 15th and 30th
 - Dry cleaners every other Monday
 - Therapy M/W/F
- Use direct deposit and automatic bill pay for as many items as possible
- Use apps such as GoodBudget to keep track of expenses



- Social Situations
 - Think ahead about situations which might result in poor judgment
 - As a caregiver, give supportive and realistic feedback when you observe the inappropriate behavior
 - Provide clear expectations before you get there
 - Plan and rehears social interactions so they will be predictable and consistent.
 - Establish a verbal and non-verbal cueing system



Conclusions

- A client / patient centered approach is central to addressing the individualized and complex nature of TBI
- Top down assessment facilitates therapeutic relationship building and a contextualized treatment approach
- Several remedial and compensatory intervention approaches exist to facilitate progression towards established goals.



Thank You





Questions?

- Steven Wheeler, PhD, OTR/L, FAOTA, CBIS
- swheeler@hsc.wvu.edu

- Janice Osborne-Dowdy, MS, CCC/SLP, CBIS
- Janice.Dowdy@neurorestorative.com



References

- Bogner J, Dijkers M, Hade E, Beaulieu C, Montgomery E, Giuffrida C, Timpson M, Peng J, Gilchrist K, Lash A, Hammond F, Horn SD, Corrigan J. (2019). Contextualized treatment in traumatic brain injury inpatient rehabilitation: effects on outcomes during the first year after discharge. *Arch Phys Med Rehabil*. 100(10):1810-1817. PMID: 30716280
- Bombardier, C., Fann, J., Ludman, E., Vannoy, S., Dyer, J., & Temkin, N, (2017). The relations of cognitive, behavioral, and physical activity variables to depression severity in traumatic brain injury: Reanalysis of data from a randomized controlled trial. *Journal of Head Trauma Rehabilitation*, 32(5), 343-353. doi: 10.1097/HTR.000000000000000288
- Hanks, R., Rapport, L., Wertheimer, J., & Koviak, C. (2012). Randomized controlled trial of peer mentoring for individuals with TBI and their significant others. *Arch Phys Med Rehabil*. 93: 1297-1304.
- Kleim J, Jones T. (2008). Principles of experience dependent neural plasticity: implications for rehabilitation after brain injury. *J Speech Lang Hear Res.* 51(1):S225-39. PMID: 18230848.
- O'Neill-Pirozzi, Kennedy, & Sohlberg, M. (2016). Evidence-based practice for the use of internal strategies as a memory compensation technique after traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 31(4), E1-E11. doi: 10.1097/HTR.00000000000181
- Wheeler S, Lane S, McMahon B. Community participation and life satisfaction following intensive, community-based rehabilitation using a life skills training approach. OTJR: Occupation, Participation and Health. 2007; 27(1), 13-22. doi:https://doi.org/10.1177/153944920702700103
- Winstein, C. & Kay, D. (2015). Translating the science into practice: Shaping rehabilitation practice to enhance recovery after brain damage. *Prog Brain Res*. 218:331-360.